

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

**THIS PAGE BLANK (USPTO)**

09/645594

#12



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 31/70, 31/715, 31/725, 31/73, 31/735, 31/72</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/17898</b> <b>(43) International Publication Date:</b> 6 July 1995 (06.07.95)
<b>(21) International Application Number:</b> PCT/CA94/00714 <b>(22) International Filing Date:</b> 29 December 1994 (29.12.94) <b>(30) Priority Data:</b> 08/175,956 30 December 1993 (30.12.93) US 08/317,228 3 October 1994 (03.10.94) US <b>(71) Applicants:</b> NOVADEX PHARMACEUTICALS LIMITED [CA/CA]; 14125 30th Avenue, South Surrey, British Columbia V4P 2J4 (CA). UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; 2075 Wesbrook Mall, Vancouver, British Columbia V6T 1W5 (CA). <b>(72) Inventors:</b> SPEERT, David, P.; 4616 West 2nd Avenue, Vancouver, British Columbia V6R 1L1 (CA). USHER, Thomas, C.; Box N7525, Nassau (BS). <b>(74) Agent:</b> DAUB, Sally, J.; Smart & Biggar, Suite 2300, 439 University Avenue, Toronto, Ontario M5G 1Y8 (CA).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> A METHOD OF PREVENTING OR REDUCING THE RISK OF INFECTION BY BACTERIAL PATHOGENS UTILIZING SIMPLE AND CONJUGATED DEXTRANS  <b>(57) Abstract</b>  Methods for reducing the risk of or preventing infections by bacterial pathogens <i>in vivo</i> . In particular, a method for reducing the risk of <i>P. aeruginosa</i> infection <i>in vivo</i> in compromised hosts such as cystic fibrosis patients. The methods involve the use of dextran or dextran sulphate as the active ingredient.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

A METHOD OF PREVENTING OR REDUCING THE RISK OF INFECTION BY  
BACTERIAL PATHOGENS UTILIZING SIMPLE AND CONJUGATED DEXTRANS

5 CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a continuation-in-part of Application Serial No. 08/175,956 filed December 30, 1993.

FIELD OF THE INVENTION

10 The present invention relates to a novel therapy for reducing the risk of acquiring bacterial infections by administering a pharmaceutical composition having simple or conjugated dextrans as active ingredient.

15 BACKGROUND OF THE INVENTION

Humans are susceptible to infection by a variety of pathogenic bacteria. Such infections pose a significant health risk for the human population in general. The human respiratory tract serves as a major portal for the  
20 opportunistic entry of bacterial pathogens into the human body. Once established and having multiplied within the respiratory tract, a bacterial pathogen may have severe localised effects or may further affect other vital bodily functions.

25 Pathogen entry into the respiratory tract can be reduced by the activity of the ciliary and mucus producing cells lining the respiratory tract. Individuals whose normal biological defence mechanisms have been compromised suffer  
30 from greater susceptibility to opportunistic infection by bacterial pathogens.

Typical examples of human diseases contracted due to the entry of bacterial pathogens into the respiratory system are bronchitis and pneumonia. Numerous bacterial species are known to be the causative agents of both of these diseases.

5 For example, Streptococcus pneumoniae is a common causative agent of pneumonia while bronchitis may be caused by numerous species of Streptococcus or Staphylococcus. For each of these disease examples, establishment of the bacterial pathogen within the respiratory system is facilitated when either the  
10 normal ciliary or mucus producing functions of the respective cells is inhibited. As well, contraction of either of these diseases often leads to the further reduction in the functional abilities of the ciliary and mucus producing cells.

15 There is, therefore, a need to provide a prophylactic therapy to reduce the risk of, or for the prevention of, infection of the respiratory system by bacterial pathogens.

20 A further example of the debilitating and often lethal effects due to infection by a bacterial pathogen is evident in persons suffering from cystic fibrosis. Cystic fibrosis ("CF") is a congenital metabolic disorder which is common among Caucasians, affecting approximately 1 in 2,000  
25 newborns. The mode of inheritance is generally autosomal recessive, suggesting that about 5% of the normal population carries the defective gene. CF manifests itself in abnormal secretions of the exocrine glands. Excessive viscid mucus

causes obstruction of passageways including pancreatic and bile ducts, the intestines, and bronchi.

Pseudomonas aeruginosa is the predominant respiratory tract pathogen in patients with cystic fibrosis. Once CF patients acquire a P. aeruginosa infection, the infection is rarely, if ever, eradicated and a progressive pulmonary deterioration is initiated, ultimately leading to death. In one study, chronic colonization established in the first five years of life was associated with a 20% survival to 16 years of age, whereas 95% of the patients who remained uncolonized in the first five years of life survived to 16 years of age. Although the extraordinary predisposition of CF patients to colonization and infection with P. aeruginosa has been recognized for many years, a satisfactory explanation for this phenomenon remains elusive.

Presently, there are currently no prophylactic measures or treatments that have resulted in the complete eradication of P. aeruginosa infections in CF patients.

CF patients with established P. aeruginosa infections have been treated by antimicrobial therapy using antibiotics in several prior art therapeutic protocols. However, the complications that have been observed in antibiotic therapy include the following. First, patients with CF dispose of antimicrobial agents more rapidly than do normal individuals, a phenomenon that mandates therapy with

higher doses than those normally recommended. Second, strains of P. aeruginosa dissociate into multiple phenotypic forms and often with different antimicrobial susceptibility patterns. Third, since the infection is chronic and the infecting strains of P. aeruginosa are rarely eradicated, resistance to multiple antimicrobial agents frequently develops. Fourth, therapeutic levels of antimicrobial agents in sputum are difficult to achieve because of poor penetration and inactivation. Fifth, the mucoid exopolysaccharide of mucoid strains appears to present a barrier to penetration of some antibiotics. Finally, allergy to certain antibiotics renders therapy with antibiotics difficult in some patients.

Further, anti-inflammatory agents have been used in the therapy of established P. aeruginosa infections in CF patients, as it has been postulated that host-mediated inflammation may be responsible for a large part of the pulmonary damage in the CF lung. Efforts have been made to dampen the inflammatory response in the CF lung by use of a systemic steroidal anti-inflammatory therapy using prednisone. Unfortunately, prednisone therapy carries substantial risks including growth retardation, glucose intolerance and development of cataracts. Preliminary studies are underway to test use of non-steroidal anti-inflammatory agents.

The manner in which P. aeruginosa establishes infection in CF patients has been studied. It is understood that a defect in CF host defences against infection must exist



a priori, otherwise infection with P. aeruginosa, an opportunistic pathogen, would not be possible. Despite extensive investigations, a unifying explanation for the peculiar predisposition of CF patients to infection with P. aeruginosa has not been forthcoming.

A range of subtle immunological abnormalities, both primary and secondary, in patients with CF has been identified and incorporated into a feasible pathophysiological scheme. The first of such abnormalities has been identified as the tendency of P. aeruginosa to adhere to buccal and respiratory epithelial cells in patients with CF.

P. aeruginosa is present in the upper and lower respiratory tracts of most older patients with CF, but it is rarely recovered from healthy individuals. The propensity of P. aeruginosa to colonize respiratory epithelia of patients with CF has been attributed to the pathogen's enhanced adhesion to buccal epithelial cells in patients with CF. It has been determined that fibronectin, a glycoprotein which normally coats buccal epithelial cells and prevents adhesion of Gram-negative bacteria, is depleted from the surface of buccal epithelial cells in patients with CF. This provides a possible explanation of the tendency of P. aeruginosa to colonize the upper respiratory tract of patients with CF.

Colonization appears to be established in the upper respiratory tract where nonmucoid P. aeruginosa adhere to

buccal epithelial cells in patients with CF, better than to the same cells in normal individuals. Contamination of the lower airways probably occurs due to spontaneous aspiration of respiratory secretions. Since there is defective mucociliary clearance in patients with CF, such aspirated secretions are not cleared and remain in the lower airways. Once P. aeruginosa reaches the lower airways, phagocytosis and killing by macrophages does not occur and a chronic state of infection is established. In this niche, the bacteria are exposed to a new environment which apparently provides the appropriate stimulus for a switch from the nonmucoid to the mucoid phenotype. These mucoid variants may have a survival advantage over nonmucoid strains by virtue of their resistance to phagocytosis and their adhesion to tracheobronchial mucin. Once chronic infection is established, the bacterial density increases dramatically in concert with a down-regulation of toxin production.

As a result of the considerable difficulty and lack of success in treating patients already infected by P. aeruginosa, there is a need to provide a prophylactic therapy to reduce the risk of, or for the prevention of, P. aeruginosa infection, most importantly in patients with CF. Further, there is a need to provide a therapy to reduce the risk of further such infection during the treatment of a patient who has already been infected.

The present invention provides a method for reducing the risk of or preventing the establishment of infection of the respiratory system by bacterial pathogens by treating individuals with a pharmaceutical composition having as an active ingredient either simple or conjugated polysaccharides. For example, treatment of the buccal or respiratory epithelial cells in a human with either simple or conjugated polysaccharides, administered through a variety of mechanisms, can achieve such a result. Use of the method of the present invention may reduce the risk of colonization of the respiratory tract by a bacterial pathogen. By inhibiting the colonization of the bacterial pathogen, the method of the present invention inhibits the establishment of a diseased state in the individual. As well, the present invention may be used for the treatment of individuals already infected by a bacterial pathogen.

Known references show pharmaceutical compositions comprising polysaccharides or their conjugates, including dextran and dextran sulphate as an active ingredient. Of such references, one has shown that such compositions may be effective in treating against infection of human cells by retroviruses. Ueno et al. ("Ueno"), in U.S. Pat. No. 5,100,879, discloses a method for the disinfection of retroviruses by the application of dextran sulphate in concentrations ranging from 0.01 to 1.0  $\mu\text{g/ml}$  to human T-lymphocytes. The molecular weight of the dextran sulphate may range from 2,000 to 10,000 and have a minimum 5% sulphur

content. Treatment of T-cells with dextran alone did not prevent T-cell infection by HTLV-III, a causative agent of AIDS. However, in view of the significant differences between the mechanism of infection of T-cells by retroviruses and infection of the respiratory system by bacterial pathogens, Ueno does not teach or suggest a treatment which would be expected to prevent or reduce the risk of respiratory system infection by bacterial pathogens.

Another reference shows a method of treatment of certain bacterial infections using extremely low concentrations of dextran sulphate. Krivan et al. ("Krivan"), in U.S. Pat. No. 5,089,479, discloses a pharmaceutical composition comprised of either sulphotides, dextran sulphate, sialyloligosaccharides or a mixture thereof, coupled with either a water insoluble or water soluble carrier. The composition is used in a method of removing the mycoplasma pathogen from patients infected with either Mycoplasma hominus or Mycoplasma pneumoniae. Also disclosed is a method for detecting mycoplasma pathogens from the said groups through the use of an enzyme-dextran sulphate conjugate.

The use of dextran sulphate at the concentrations disclosed in the present invention for reducing the risk of bacterial infection of the respiratory system is not taught or suggested by Krivan as the concentrations of dextran sulphate utilized by Krivan are approximately 1000 times lower than those disclosed herein. Further, Krivan teaches that dextran

has no effect on the treatment of Mycoplasma infections and therefore teaches away from the invention disclosed presently.

Methods for treating patients having CF are known.

5 For example, Edwards et al. ("Edwards"), in U.S. Pat. No. 4,866,072, discloses a method for the treatment of CF by the direct administration of the active ingredient nedocromil sodium, in an aqueous solution in aerosol or oral format, to a patient's lung. The active ingredient may range in  
10 concentration from 0.1% to 2.0% w/w for aqueous solutions and from 0.5% to 5.0% w/w for pressurized aerosols. Edwards does not contemplate a method for treatment of the respiratory system against bacterial infection.

15 SUMMARY OF THE INVENTION

According to an aspect of the present invention, there is provided a method for reducing the risk of respiratory system infection by bacterial pathogens which comprises administering to a patient in need of such treatment  
20 a pharmaceutical composition comprising an effective amount of a simple or conjugated polysaccharide in admixture with a pharmaceutically acceptable diluent or carrier.

According to another aspect of the present  
25 invention, there is provided a method for reducing adhesion of bacterial pathogens to buccal or respiratory epithelial cells *in vivo* which comprises administering to a patient in need of such treatment a pharmaceutical composition comprising an

effective amount of a simple or conjugated polysaccharide in admixture with a pharmaceutically acceptable diluent or carrier.

5           According to yet another aspect of the present invention, there is provided a method reducing the risk of respiratory tract infection by P. aeruginosa in a patient having cystic fibrosis which comprises administering to said patient a pharmaceutical composition comprising an effective  
10   amount of dextran sulphate or dextran in admixture with a pharmaceutically acceptable diluent or carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

15           Figure 1 shows the effect of dextran sulphate on the adhesion of P. aeruginosa to buccal epithelial cells.

          Figure 2 shows the effect of dextran on the ability of P. aeruginosa to grow in culture.

20           Figure 3 shows the effect of low molecular weight dextran on the growth of P. aeruginosa in culture.

          Figure 4 shows the effect of dextran upon the ability of P. aeruginosa isolated from CF patients to adhere  
25   to human respiratory epithelial cells.

Figure 5 shows the ability of dextran of various molecular weights to inhibit the adhesion of P. aeruginosa PAK to human respiratory epithelial cells.

5 Figure 6 shows the ability of Dextran 4000 at various concentrations to inhibit the adhesion of P. aeruginosa PAK to human respiratory epithelial cells.

10 Figures 7 and 8 show the effect of Dextran 4000 to inhibit the adhesion of Gram positive bacteria to human respiratory epithelial cells.

#### DETAILED DESCRIPTION OF THE INVENTION

15 The inventors have found that dextran or dextran sulphate may be used to reduce the risk of or prevent the adhesion of bacterial pathogens to human respiratory epithelial cells in vitro. From this it can be soundly predicted that polysaccharides such as dextran or dextran sulphate will be effective in reducing the risk of or  
20 preventing bacterial pathogens infections in compromised hosts (such as patients with CF) by blocking adhesion (and therefore colonization) to the airway epithelial cells. As well, polysaccharides such as dextran or dextran sulphate would be  
25 effective in treating patients already infected with P. aeruginosa by reducing the risk of or preventing further infection of such hosts.

Dextran sulphate and its known uses will be recognized by persons skilled in the art. For example, the salts of dextran sulphate, particularly its sodium salt, have been found to be useful as an anti-coagulant, as an  
5 anti-lipemic agent and an anti-ulcer agent.

Dextran sulphate is a sulphonated polysaccharide having a wide range of molecular weights. Based upon their studies, the inventors recommend that dextran sulphate having  
10 an average molecular weight in the range of 8,000 - 1,000,000 would be most effective in reducing adhesion of bacterial pathogens to buccal or respiratory epithelial cells.

Experimental results supporting the finding that  
15 dextran sulphate reduces the adhesion of bacterial pathogens to buccal epithelial cells are detailed below.

Dextran and its known uses will also be recognized by persons skilled in the art. For example, dextran is known  
20 to be useful for the expansion of blood plasma. As well, dextran can act as a blood flow adjuvant due to its ability to prevent erythrocyte aggregation.

Hydrogenated dextran, commonly referred to as simply  
25 dextran, is a polymer of glucose, possessing a wide range of molecular weights. Dextran is commercially produced by growing bacterium (Leuconostoc mesenteroides) upon a sucrose substrate to yield high molecular weight dextrans.



Depolymerization and fractionation of the high molecular weight dextrans will yield lower weight dextrans. Based upon their studies, the inventors recommend that dextran having an average molecular weight in the range of 3,000 - 1,000,000 would be most effective in reducing adhesion of bacterial pathogens to human respiratory epithelial cells.

Experimental results supporting the finding that dextran reduces the adhesion of bacterial pathogens to buccal epithelial cells are also detailed below.

#### MATERIALS AND METHODS

##### Dextran Sulphate

Fresh buccal epithelial cells were obtained from the mouths of two healthy individuals by gently swabbing with alginate swabs. The epithelial cells were suspended in phosphate buffered saline and washed. The cells were divided in tubes into 3 groups for (i) a control; (ii) addition of P. aeruginosa and (iii) addition of P. aeruginosa and dextran sulphate. P. aeruginosa was added to one of the tubes containing the fresh epithelial cells. P. aeruginosa and dextran sulphate were added to another tube containing the fresh epithelial cells. The dextran sulphate used is sold under the trademark Usherdex 8 and is a hydrogenated form of dextran sulphate having an average molecular weight of around 8,000. The dextran sulphate solution was 10 mmol strength. All of the tubes were incubated for 60 minutes. At the end of the incubation, the cells were washed and deposited by

cytopsin on glass slides. The slides were allowed to dry overnight and then stained with Giemsa stain and the bacteria associated with each epithelial cell was determined by visual inspection. Experimental results are demonstrated in

5 Figure 1.

Referring to Figure 1, three different conditions are depicted by separate bars on the graph of Figure 1. The three sets of bars represent the following (from left to

10 right):

(a) the number of bacteria which are normally resident on the epithelial cells from the mouth (ie. no dextran sulphate or P. aeruginosa were added);

15

(b) the results of the experiment in which P. aeruginosa was added; and

20

(c) the results when dextran sulphate (10 mmol) was added along with the P. aeruginosa to the epithelial cells.

25

The adhesion of P. aeruginosa to buccal epithelial cells is shown in Figure 1 by comparing the second and first set of bars. The addition of the bacteria resulted in increased incidence of bacteria per epithelial cell over the number found normally in the mouth. Comparing the third and second set of bars indicates that the treatment of the

epithelial cells with dextran sulphate resulted in a reduction in the adhesion of the bacteria to the cells.

Up to 20 mmol (160 mg/ml) dextran sulphate has been used with similar results. As demonstrated in Figure 1, it has been found that a concentration of 10 mmol (80 mg/ml) is highly effective at interfering with adhesion of P. aeruginosa to epithelial cells. Concentrations in the range of 1-50 mmol would be useful for the purpose described herein. It is believed that dextran sulphate concentrations in the range of 10-20 mmol would be most useful in reducing the risk of such P. aeruginosa infections.

It is believed that dextran sulphate with a molecular weight in the range of 8,000 to 1,000,000 would be effective in a treatment to reduce the risk of bacterial infections.

In vivo, the dextran sulphate would preferably be administered topically. Dextran sulphate as the active ingredient may be administered topically by gargle, swish and swallow, oral lozenge, aerosol or other such manner. Dextran sulphate and suitable carrier compositions will be understood by those skilled in the art. Preferably, dextran sulphate is delivered to the site of potential adhesion to buccal epithelial cells in concentrations as mentioned above.

Although the foregoing description suggests specific manners of topical administration of dextran sulphate to buccal epithelial cells, it will be understood by persons skilled in the art that other methods of administration are possible.

As a result of its anti-coagulant properties, dextran sulphate may cause some undesired side effects in use. The dextran sulphate used in the present experiment was in its hydrogenated form. Hydrogenation reduces the anti-coagulant effect of the dextran sulphate and is therefore the preferred form as an active ingredient in methods and compositions described herein. It is expected that dextran sulphate in its unhydrogenated form would also be effective for use in a treatment to reduce the risk of such infections.

Preferably, a patient would be treated 2-3 times daily with dextran sulphate solution. Greater and lesser frequencies may be effective depending upon the individual patient's requirements. It is believed that at least daily treatment would best reduce the risk of P. aeruginosa infection.

It is believed that dextran sulphate may also be effective in assisting with the treatment of patients already infected by P. aeruginosa by reducing the risk of re-colonization of the patient.

Further, it will be appreciated by persons skilled in the art that other, non-topical methods of treatment using dextran sulphate can be successful in reducing the risk of or preventing infection by P. aeruginosa in compromised hosts.

5 It is known that dextran sulphate of average molecular weight of 8,000 is absorbed by the body and circulated systemically. The dextran sulphate may therefore be delivered to sites of P. aeruginosa adhesion and secreted to have effects equivalent to that of the topically applied form. Such methods include  
10 systemic therapy, i.e. the ingestion per os of dextran sulphate in tablet form.

The foregoing detailed description and examples are for purposes of illustration only, and are not intended as  
15 limiting the scope of the invention. Persons skilled in the art will appreciate the nature and scope of the invention including its practical application.

#### Dextran

20 The methodology described below details the experiments performed to determine the effect of dextran on the growth of bacterial cells.

25 Bacteria were inoculated into minimal medium (BM2) containing L-methionine and glucose. Alternatively, bacteria were inoculated into LB broth. The bacterial strain inoculated into the LB broth was P. aeruginosa PAK, while the bacterial strain inoculated into the BM2 broth was

P. aeruginosa P1. Bacterial growth was monitored by reading the culture absorbency at 600 nm. To test the effect of dextran upon bacterial growth, 8.6 mmol dextran 6,000 (a hydrogenated form of dextran sold under the trademark "Usherdex 6000") was added to the BM2 medium, while 40 mg/ml of either dextran 4,000 (a hydrogenated form of dextran sold under the trademark "Usherdex 4000") or dextran 40,000 was added to the LB broth. These results are shown in Figures 2 and 3. In all cases, dextran solutions were adjusted to a pH of between 7.0-7.2 by the addition of 1N NaOH.

Referring to Figure 2, the open circles represent the growth rate of P. aeruginosa, as measured by a spectrophotometer absorbance reading at 600 nm of the liquid bacterial culture, without the addition of dextran to the bacterial growth medium. The darkened circles represent the growth rate of P. aeruginosa, as measured by a spectrophotometer absorbance reading at 600 nm of the liquid bacterial culture, after the addition of dextran the bacterial growth medium. Comparison of the two growth rates indicates that dextran does not provide a rich growth medium for P. aeruginosa in vitro.

Referring to Figure 3, upper diagram, the open circles represent the number of colony forming units present in the liquid culture of P. aeruginosa without the addition of dextran the bacterial growth medium. The darkened circles represent the number of colony forming units present in the

liquid culture of P. aeruginosa when 40 mg/ml of dextran 4,000 is added to the bacterial growth medium. The open triangles represent the number of colony forming units present in the liquid culture of P. aeruginosa when 40 mg/ml of dextran 40,000 is added to the bacterial growth medium.

Comparison of the number of colony forming units present for any given time when P. aeruginosa is cultured in the absence of dextran to the number of colony forming units present for any given time when P. aeruginosa is cultured in the presence of either dextran 4,000 or dextran 40,000 indicates that neither dextran 4,000 or dextran 40,000 provide a rich growth medium for P. aeruginosa bacterial cells.

Comparison of the number of colony forming units present for any given time when P. aeruginosa is cultured in the presence of dextran 4,000 as opposed to dextran 40,000 indicates that dextran 4,000 has a greater ability than dextran 40,000 to reduce the number of P. aeruginosa bacterial cells which will grow in culture.

Referring to Figure 3, lower diagram, the open circles represent the number of P. aeruginosa cells present in the liquid culture, as measured by a spectrophotometer absorbance reading at 600 nm of the liquid bacterial culture, present in the liquid culture of P. aeruginosa without the addition of dextran to the bacterial growth medium. The darkened circles represent the number of P. aeruginosa cells

present in the liquid culture, as measured by a spectrophotometer absorbance reading at 600 nm of the liquid bacterial culture, when 40 mg/ml of dextran 4,000 is added to the bacterial growth medium. The open triangles represent the number of P. aeruginosa cells present in the liquid culture, as measured by a spectrophotometer absorbance reading at 600 nm of the liquid bacterial culture, when 40 mg/ml of dextran 40,000 is added to the bacterial growth medium.

Comparison of the number of P. aeruginosa cells present in the liquid culture for any given time when P. aeruginosa is cultured in the absence of dextran to the number of P. aeruginosa cells present in the liquid culture for any given time when P. aeruginosa is cultured in the presence of either dextran 4,000 or dextran 40,000, indicates that neither dextran 4,000 and dextran 40,000 provide a rich growth medium for P. aeruginosa bacterial cells.

Comparison of the number of P. aeruginosa cells present in the liquid culture, up to 4.5 hours after the initiation of bacterial growth in the liquid media, when P. aeruginosa is cultured in the presence of dextran 4,000 as opposed to dextran 40,000 indicates that dextran 4,000 has a greater ability than dextran 40,000 to reduce the number of P. aeruginosa bacterial cells which will grow in culture.

The methodology described below was employed in experiments to determine the effects of dextran upon the



ability of bacterial cells to adhere to A549 human lung epithelial cells.

A549 human lung epithelial cells were obtained from the American Type Culture Collection (ATCC # CCL 185). The A549 cells were maintained in F-12K medium and harvested by trypsin treatment as recommended by ATCC. Cells were plated at a density of  $2 \times 10^4$  cells/well on a coverslip placed in a 24-well tissue culture tray (Becton Dickson, Lincoln Park, NJ). After 2 hours or 24 hours, cells were washed with PBS (pH 7.0, from Oxoid, Nepean, Ontario, Canada) at 37°C. These were overlaid with 500  $\mu$ l of Binding Assay Buffer (BAB; 138 mmol NaCl, 8.1 mmol  $\text{Na}_2\text{HPO}_4$ , 1.5 mmol  $\text{KH}_2\text{PO}_4$ , 2.7 mmol KCl, 1.0 mmol  $\text{MgCl}_2$ , 0.25 mmol  $\text{CaCl}_2$  and 0.001% phenol red, pH 7.2; Sigma Chemical Company, St. Louis, MO). Cells were incubated for 30 minutes at 37°C in BAB or 10 mmol dextran made in BAB. Subsequently, 25  $\mu$ l of bacteria (grown in L-broth to  $\text{OD}_{600}=0.5$ ) were added and the entire epithelial cell - bacterial cell mixture was re-incubated for another 30 minutes. The coverslips were then washed four times with PBS, fixed in methanol for 15 minutes, mounted on glass microscope slides, dried for 60 minutes at 37°C and stained for 10 to 15 minutes in Giemsa. Bound bacteria were assessed microscopically by surveying 60 epithelial cells per co-incubation treatment. The average number of bacteria/cell was calculated.

Referring to Figure 4, a comparison of the open scale-bars with the closed scale-bars in each column indicates

that the addition of the dextran to the culture media caused a reduction of the adhesion of P. aeruginosa bacterial cells to the epithelial cells growing in vitro. The results indicate that the reduction in the adhesion of the bacteria was independent of the bacterial isolate. Each bacterial isolate was obtained from a different CF patient (except for strain PAK).

Up to 20 mmol dextran has been used with similar results. A concentration of 10 mmol for dextran 6,000 has been shown to be highly effective at interfering with the adhesion of P. aeruginosa to the epithelial cells. Concentrations in the range of 1 mmol to 50 mmol would be useful for the purpose described herein.

Referring to Figure 5, the scale-bars represent the number of bacteria which were observed to adhere to the epithelial cells after 30 minutes of co-incubation of the bacterial and epithelial cells. The addition of dextran to the growth media resulted in a decrease in adherence of P. aeruginosa to the epithelial cells. On a comparison of equal weight to volume ratios, dextran 3,000 was most effective in reducing the adhesion of the bacteria to the epithelial cells. The use of dextran with a molecular weight in the range of 3,000 to 1,000,000 would be effective in interfering with the adhesion of P. aeruginosa to epithelial cells.

Referring to Figure 6, the results indicate that the addition of dextran 4,000 in concentrations of 5 mmol or greater is highly effective in reducing the adhesion of P. aeruginosa to epithelial cells.

5

Referring to Figure 7, the darkened scale-bars represent the number of bacteria observed to be adhering to the individual epithelial cells after 30 minutes of co-incubation of the epithelial cells with the bacterium in the absence of dextran. The open scale-bars represent the number of bacteria observed to be adhering to the individual epithelial cells after 30 minutes of co-incubation of the epithelial cells with the bacterium in the presence of 10 mmol dextran 4,000.

15

For the first and second set of scale-bars, a comparison of the darkened and open scale-bars within each set indicates that the addition of the 10 mmol dextran 4,000 to the culture media during the co-incubation stage is highly effective in reducing the adherence of Gram positive bacteria (Enterococcus and S. aureus respectively) to the epithelial cells. The third set of scale-bars indicates that Streptococcus epidermidis, a Gram positive bacteria, did not bind to the epithelial cells even in the absence of dextran.

25

Referring to Figure 8, the darkened scale-bars represent the number of bacteria observed to be adhering to the individual epithelial cells after 30 minutes of

co-incubation of the epithelial cells with the bacterium in the absence of dextran. The open scale-bars represent the number of bacteria observed to be adhering to the individual epithelial cells after 30 minutes of co-incubation of the epithelial cells with the bacterium in the presence of 10 mmol dextran 4,000.

For each strain of Streptococcus pneumoniae tested, a comparison of the darkened and open scale-bars within each set indicates that the addition of the 10 mmol dextran 4,000 to the culture media during the co-incubation stage is highly effective in reducing the adherence of Gram positive bacteria to the epithelial cells.

In vivo, dextran would preferably be administered topically. Dextran as the active ingredient may be administered topically by gargle, swish and swallow, oral lozenge, aerosol or other such manner. Dextran and suitable carrier compositions will be understood by those skilled in the art. Preferably, dextran is delivered to the site of potential adhesion to buccal or respiratory epithelial cells in concentrations as mentioned above.

Although the foregoing description suggests specific manners of topical administration of dextran to buccal or respiratory epithelial cells, it will be understood by persons skilled in the art that other methods of administration are possible.

The dextran used in the present experiments was in its hydrogenated form. It is expected that dextran in its unhydrogenated form would also be effective for use in a treatment to reduce the risk of the type of bacterial infections described herein.

Preferably, a patient would be treated 2-3 times daily with dextran solution. Greater and lesser frequencies may be effective depending upon the individual patient's requirements. It is believed that at least daily treatment would best reduce the risk of P. aeruginosa infection.

It is believed that dextran may also be effective in assisting with the treatment of patients already infected by P. aeruginosa by reducing the risk of re-colonization of the patient.

Further, it will be appreciated by persons skilled in the art that other, non-topical methods of treatment using dextran can be successful in reducing the risk of or preventing infection by P. aeruginosa in compromised hosts. It is known that dextran of average molecular weight is absorbed by the body and circulated systemically. The dextran may therefore be delivered to sites of P. aeruginosa adhesion and secreted to have effects equivalent to that of the topically applied form. Such methods include systemic therapy, i.e. the ingestion per os of dextran in tablet form.

It may be soundly predicted on the basis of the foregoing results that sulphonated polysaccharides selected from the group consisting of cellulose sulphate, chondroitin sulphate, sucrose sulphate and heparin would be effective for the purposed of the method disclosed herein. Further, other polysaccharides selected from the group consisting of glycosaminoglycan, hyaluronic acid, chondroitin sulphate, dermatin sulphate, heparin sulphate, keratin sulphate and keratin would also be expected to be effective.

10

The foregoing detailed description and examples are for purposes of illustration only, and are not intended as limiting the scope of the invention. Persons skilled in the art will appreciate the nature and scope of the invention including its practical application.

15

## WE CLAIM:

1. A method for reducing the risk of respiratory system infection by bacterial pathogens which comprises administering to a patient in need of such treatment a pharmaceutical composition comprising an effective amount of a simple or conjugated polysaccharide in admixture with a pharmaceutically acceptable diluent or carrier.

2. The method of claim 1 wherein said simple or conjugated polysaccharide is a sulphonated polysaccharide.

3. The method of claim 2 wherein said sulphonated polysaccharide is hydrogenated.

4. The method of claim 2 wherein said sulphonated polysaccharide is selected from the group consisting of cellulose sulphate, chondroitin sulphate, sucrose sulphate, heparin or dextran sulphate.

5. The method of claim 4 wherein said sulphonated polysaccharide is dextran sulphate.

6. The method of claim 5 wherein said dextran sulphate is hydrogenated.

7. The method of claim 5 wherein said carrier is a liquid, an oral lozenge or an aerosol.

8. The method of claim 7 wherein said dextran sulphate is administered in liquid form by gargle or swish and swallow.

9. The method of claim 7 wherein said dextran sulphate is present in a final concentration of about 1-50 mmol.

10. The method of claim 9 wherein said dextran sulphate is present in a final concentration of about 10-20 mmol.

11. The method of claim 10 wherein said dextran sulphate is present in a final concentration of about 10 mmol.

12. The method of claim 7 wherein said dextran sulphate concentration at a potential site of bacterial pathogen adhesion is about 10-20 mmol.

13. The method of claim 12 wherein said dextran sulphate concentration at a potential site of bacterial pathogen adhesion is about 10 mmol.

14. The method of claim 7 wherein said dextran sulphate is hydrogenated.

15. The method of claim 7 wherein said bacterial pathogen is Pseudomonas aeruginosa, Streptococcus epidermidis, Staphylococcus aureus, Streptococcus pneumoniae or Enterococcus.



16. The method of claim 7 wherein said dextran sulphate has an average molecular weight of about 8,000 - 1,000,000.

17. The method of claim 16 wherein said dextran sulphate has an average molecular weight of 8,000.

18. The method of claim 1 wherein said simple or conjugated polysaccharide is hydrogenated dextran.

19. The method of claim 18 wherein said carrier is a liquid, an oral lozenge or an aerosol.

20. The method of claim 19 wherein said dextran is administered in liquid form by gargle or swish and swallow.

21. The method of claim 19 wherein said dextran is present in a final concentration of about 1-50 mmol.

22. The method of claim 21 wherein said dextran is present in a final concentration of about 10-20 mmol.

23. The method of claim 22 wherein said dextran is present in a final concentration of about 10 mmol.

24. The method of claim 19 wherein said dextran concentration at a potential site of bacterial pathogen adhesion is about 10-20 mmol.

25. The method of claim 24 wherein said dextran concentration at a potential site of bacterial pathogen adhesion is about 10 mmol.

5 26. The method of claim 19 wherein said bacterial pathogen is Pseudomonas aeruginosa, Streptococcus epidermidis, Staphylococcus aureus, Streptococcus pneumoniae or Enterococcus.

10 27. The method of claim 18 wherein said dextran has an average molecular weight of about 3,000 - 1,000,000.

28. The method of claim 26 wherein said dextran has an average molecular weight of about 4,000 - 6,000.

15 29. The method of claim 27 wherein dextran has an average molecular weight of about 4,000.

20 30. The method of claim 1 where said simple or conjugated polysaccharide is selected from the group consisting of unhydrogenated dextran, glycosaminoglycan, hyaluronic acid, chondroitin sulphate, dermatin sulphate, heparin sulphate, heparin, keratin sulphate or keratin.

25 31. A method for reducing adhesion of bacterial pathogens to buccal or respiratory epithelial cells in vivo which comprises administering to a patient in need of such treatment a pharmaceutical composition comprising an effective


amount of a simple or conjugated polysaccharide in admixture with a pharmaceutically acceptable diluent or carrier.


32. The method of claim 31 wherein said bacterial pathogens are Pseudomonas aeruginosa, Streptococcus epidermidis, Staphylococcus aureus, Streptococcus pneumoniae or Enterococcus.


33. The method of claim 32 wherein said simple or conjugated polysaccharide is selected from the group consisting of dextran sulphate or dextran.


34. The method of claim 33 wherein said bacterial pathogen is Pseudomonas aeruginosa and said patient has cystic fibrosis.


35. A method reducing the risk of respiratory tract infection by Pseudomonas aeruginosa in a patient having cystic fibrosis which comprises administering to said patient a pharmaceutical composition comprising an effective amount of dextran sulphate or dextran in admixture with a pharmaceutically acceptable diluent or carrier.




**FIG. 2**





FIG. 4

FIG. 5



FIG. 6

FIG. 7

**FIG. 8**

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 94/00714

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/70 A61K31/715 A61K31/725 A61K31/73 A61K31/735  
A61K31/72

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 126 043 (SVENSKA SOCKERFABRIKS AB) 21 November 1984	1,30-32
Y	* p.1, 4-25; claims 6,11; Tables I-IV * see the whole document	2-30, 33-35
X	WO,A,93 07178 (PASTEUR MERIEUX SERUMS ET VACCINS) 15 April 1993	1,30-32
Y	* claims 7,9,16,18 * see the whole document	2-29, 33-35
X	WO,A,93 24505 (ALBERTA RESEARCH COUNCIL) 9 December 1993	1-4, 30-32
Y	* claims 2,4 * see the whole document	5-29, 33-35
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier document but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 May 1995

Date of mailing of the international search report

06.06.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+ 31-70) 340-3016

Authorized officer

Uiber, P

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 94/00714

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,93 08810 (CARRINGTON LAB. INC.) 13 May 1993 * p.5, 1.5-13; p.26, 1.19; Ex. 35, 45 * ---	1-35
Y	WO,A,91 15216 (KENNEDY T.P.) 17 October 1991 * p.1, 1.17-30; claims 3,13 * ---	1-35
Y	US,A,5 217 715 (KRIVAN ET AL) 8 June 1993 see the whole document ---	1-35
A	EP,A,0 177 783 (KANTO ISHI PHARMACEUTICAL CO, LTD) 16 April 1986 see the whole document -----	1-35

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 94/00714

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-126043	21-11-84	AU-B- 566286	15-10-87
		AU-A- 2587184	27-09-84
		JP-A- 59176300	05-10-84
-----			
WO-A-9307178	15-04-93	FR-A- 2682388	16-04-93
		AU-A- 2946992	03-05-93
		EP-A- 0562107	29-09-93
		FI-A- 932626	09-06-93
		JP-T- 6506233	14-07-94
-----			
WO-A-9324505	09-12-93	CA-A- 2110707	23-12-92
		CA-A- 2110997	23-12-92
		CA-A- 2118405	12-09-93
		CA-A- 2118522	09-12-93
		WO-A- 9222301	23-12-92
		WO-A- 9222564	23-12-92
		WO-A- 9222662	23-12-92
		EP-A- 0591254	13-04-94
		EP-A- 0591256	13-04-94
		EP-A- 0588852	30-03-94
		EP-A- 0650492	03-05-95
		EP-A- 0643718	22-03-95
		JP-T- 7502011	02-03-95
		JP-T- 6510746	01-12-94
		JP-T- 6510661	01-12-94
		WO-A- 9324506	09-12-93
		US-A- 5374655	20-12-94
-----			
WO-A-9308810	13-05-93	EP-A- 0611304	24-08-94
		JP-T- 7500568	19-01-95
-----			
WO-A-9115216	17-10-91	AU-A- 7681091	30-10-91
-----			
US-A-5217715	08-06-93	US-A- 5389521	14-02-95
		US-A- 5386027	31-01-95
		US-A- 5225330	06-07-93
-----			
EP-A-177783	16-04-86	JP-C- 1478542	27-01-89
		JP-A- 61068418	08-04-86
		JP-B- 63028411	08-06-88

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 94/00714

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-177783		AT-T- 111738	15-10-94
		AU-A- 4719985	20-03-86
		DE-D- 3587587	21-10-93
		DE-T- 3587587	31-03-94
		DE-D- 3587928	27-10-94
		DE-T- 3587928	26-01-95
		EP-A, B 0429430	29-05-91
		US-A- 4698332	06-10-87
-----			

**THIS PAGE BLANK (USPTO)**